



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

probability arises that they will be left high and dry on the shore, they can appreciate the fact by the growing intensity of the light, and by that peculiar flapping motion of the valves the Pectens are so remarkable for, move away into deeper water.' This theory may at once be set aside when we consider that the Pectens of the Mediterranean, where we have practically no tide at all, a state of affairs that has existed for an exceedingly long period of time, have as well developed eyes as those found on shores where tides do exist; and further, it seems hardly probable that such a complicated organ would have developed to determine for the animal whether it be out of water or not. . . . As regards the complicated organ known as the eye, I might suggest that, if this be an eye, it is one where we have no true pigmented layer in any direct relation to either the nerve or the retina. The mass of red pigment and the metallic-like tapetum would hardly answer the place of the black choroid coat so essential to the eye."

Dr. Sharp stated that at the time the above was written, he was under the impression that the organ was probably a phosphorescent organ, but he had no proof of it. At Nantucket he obtained many specimens of the animal and found that the edge of the mantle was phosphorescent, and on questioning Dr. Kite, formerly of the "Fish-Hawk," who had often seen *Pecten* dredged at night, he was informed that the phosphorescent condition of Pectens had often been observed. He thought it was not unreasonable to suppose that organs for the emission of light should be formed upon the same general principles as organs for the admission of light, hence the similarity of these organs to eyes. He further stated that this function (phosphorescence) would be of great use to the animal in obtaining its food.

On Amia and its probable Tænia.—Prof. LEIDY stated that in our market on Saturday last, three Mud-fishes, *Amia calva*, had been given to him. They came in a barrel of shad from North Carolina. One was a female about two feet long, the others male, of which the smallest was eight inches. Protruding from the vent of the latter was a little tape-worm, which, on disturbance, retreated into the rectum. Three other worms of the same kind were found in the mouth, but none were in the intestine of this or the other fishes. The worms accord with the description of the *Tænia filicolis*, infesting Sticklebacks, *Gasterosteus*, and is probably the same species. They range from $1\frac{1}{2}$ to 3 inches long, gradually widening from the delicate thread-like neck to the posterior rounded extremity where they measure from 1 to 1.5 mm. wide. The head is spheroidal, variably broader or longer, and about 0.625 mm.; with the summit slightly prominent and unarmed, and with four hemispherical, lateral bothria 0.25 mm. in diameter. Neck variable; when extended, long and narrow and usually about half the width of the head. Anterior segments,

transversely linear, about an eighth the length of the breadth; gradually becoming inverted saucer-shaped or scutellate, and about one-fourth the length of the breadth. Posterior segments more quadrate, slightly widening behind, about 0.75 mm. long and from 1. to 1.5 mm. broad. Last segment longest and rounded. Genital apertures marginal.

MARCH 30.

Mr. CHARLES MORRIS in the chair.

Twelve persons present.

Prof. J. C. Arthur was elected a correspondent.

The following was ordered to be printed :—